REPORT DOCUMENTATION PAGE

CMB No. 0704-0188

Public reporting burden for this reliection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the sollection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington releasuanters Services, Directorate for information Operations and Peports, 1215 Jefferson Calls Birdhaw, Suite 1204, Arlington, VA 12202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, EC 20503. 3. REPORT TYPE AND DATES COVERED 1. AGENCY USE ONLY (Leave blank) 2. REPORT DATE FINAL/01 OCT 92 TO 30 SEP 94 5. FUNDING NUMBERS 4. TITLE AND SUBTITLE (HBCU/MI FY92-FY93) A CLUSTER WORKSTATION FACILITY FOR RESEARCH IN COMPUTER SCIENCE (U) 6. AUTHOR(S) 4276/AS F49620-93-1-0001 Professor Thomas Mathew 8. PERFORMING ORGANIZATION 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) REPORT NUMBER University of Maryland Baltimore County APOLE-TR- .95 00/8 Baltimore MD 21228 10. SPONSORING / MONITORING 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AGENCY REPORT NUMBER AFOSR/NM F49620-93-1-0001 110 DUNCAN AVE, SUITE B115 BOLLING AFB DC 20332-0001 11. SUPPLEMENTARY NOTES FEB 23 1995 12b. DISTRIBUTION CODE 12a. DISTRIBUTION / AVAILABILITY STATEMENT UL APPROVED FOR PUBLIC RELEASE: DISTRIBUTION IS UNLIMITED 13. ABSTRACT (Maximum 200 words) The proposal addressed some statistical inference problems in the areas of multivariate calibration and meta-analysis. Multivariate calibration deals with using the statistical relationship between a response variable and an explanatory variable for statistical inference concerning an unknown value of the explanatory variable using available data. The problems that have been solved deal with the construction of confidence regions for the unknown value of the explanatory variable. Satisfactory solutions to some open problems in this area have been obtained. Meta-analysis deals with combining several independent tests concerning a common parameter. Five papers have been written based on the proposal, of which three are already published. In addition, two papers are currently under DIEG CUALITY THE STATE O preparation. 15. NUMBER OF PAGES 14. SUBJECT TERMS 16. FRICE CODE SECURITY CLASSIFICATION | 20. LIMITATION OF ABSTRACT

> Standard Form 298 (Rev. 2-89)

SAR(SAME AS REPORT):

NSN 7540-01-080-5800

SECURITY CLASSIFICATION

OF REFORT

UNCLASSIFIED

UNCLASSIFIED

OF THIS PAGE

SECURITY CLASSIFICATION

UNCLASSIFIED

OF ABSTRACT


Final Technical Report, Oct. 1992 – September 1994

Statistical Inference Problems in Some Multivariate Linear Models With Applications to Multivariate Calibration and Meta-Analysis (AFOSR F49620-93-1-0001)

Thomas Mathew

University of Maryland Baltimore County Baltimore, Maryland 21228

Air Force Office of Scientific Research 110 Duncan Avenue, Suite B115 Bolling AFB, DC 20332



19950214 007

Distribution Unlimited

Abstract: The proposal addressed some statistical inference problems in the areas of multivariate calibration and meta-analysis. Multivariate calibration deals with using the statistical relationship between a response variable and an explanatory variable for statistical inference concerning an unknown value of the explanatory variable using available data. The problems that have been solved deal with the construction of confidence regions for the unknown value of the explanatory variable. Satisfactory solutions to some open problems in this area have been obtained. Meta-analysis deals with combining information from several independent sources to arrive at an overall combined decision. The problems that have been solved in this context deal with combining several independent tests concerning a common parameter. Five papers have been written based on the proposal, of which three are already published. In addition, two papers are currently under preparation.

calibration, confidence region, meta-analysis multivariate linear model

4

UNCLASSIFIED

UNCLASSIFIED

UNCLASSIFIED

SAR

The problems formulated in the proposal were in the areas of multivariate calibration and meta-analysis, when the data can be modeled using a multivariate linear model. What follows is a very brief description of the above problems, along with a summary of the work completed and papers prepared under my AFOSR Grant (AFOSR F49620-93-1-0001). I would like to mention at the very beginning that among the work completed under the grant, I consider my work on multivariate calibration to be quite significant. I anticipate that the results I have obtained in this context will be widely used in practical problems dealing with calibration, notably in chemometrics and in spectroscopy. (In response to a request from the Program Director Dr. Jon Sjogren, I selected a paper on calibration — paper #4 below — as my best research accomplishment since October 1993).

Statistical calibration deals with procedures to determine the characteristics of an object (which are difficult to measure directly), using easily obtained measurements on related variables, by means of a statistical model that relates the characteristics of interest with the variables on which measurements can be made. My work in this area, funded by AFOSR, has been on the construction of confidence regions. The area of meta-analysis deals with quantitative procedures for combining information from several independent sources. In this context, the problems formulated in the proposal dealt with combining several independent tests. The following papers have been completed under the grant (part of the work reported in papers #1 and #2 below have been completed under my earlier AFOSR grant, AFOSR 89-0237):

- 1. (With S. Kasala) An exact confidence region in multivariate calibration. Appeared in *The Annals of Statistics*, 1994.
- 2. (With L. Zhou) Combining independent tests in multivariate linear models. Appeared in *Journal of Multivariate Analysis*, 1994.
- 3. (With A. Khuri and D. Nel) A test to determine closeness of multivariate Satterthwaite's approximation. Appeared in *Journal of Multivariate Analysis*, 1994.
- 4. (With W. Zha) A conservative confidence region in multivariate calibration. Submitted to The Annals of Statistics.
- 5. (With K. Nordström) An inequality for a measure of deviation in linear models. Submitted to *The American Statistician*.

In addition, the following papers are under preparation.

- 6. (With W. Zha). Multiple use confidence regions in multivariate calibration.
- 7. (With R. J. Kelly). Nonnegative estimation of variance components in some time

series models.

The research work in papers #1, #4 and #6 deal with the construction of confidence regions in the statistical calibration problem. The set up is as follows. Consider the $N \times p$ matrix Y of observations following the multivariate linear model

$$Y = BX + E$$

where B is a $p \times m$ matrix of unknown parameters, X is a known $m \times p$ matrix and E is a matrix of error terms. The rows of E are assumed to be independent having the multivariate normal distribution $N(0,\Sigma)$, the covariance matrix Σ being unknown. The N rows of Y represent N independent observations obtained based on known design points given by the different rows of X. If y_0 represents a future observation corresponding to an unknown design point θ , we get the model

$$\mathbf{y}_0 = B\theta + \mathbf{e}_0$$

where the error term \mathbf{e}_0 is again assumed to follow the $N(0,\Sigma)$ distribution, and is independent of E. Thus the calibration problem consists of estimating B and Σ using Y and using it, along with \mathbf{y}_0 , to obtain information about θ . We have addressed the problem of constructing confidence regions for θ . In spite of the fact that statistical calibration is a widely used procedure, a satisfactory solution has not been available for this problem, except asymptotically. The procedures that were available in the finite sample case have serious drawbacks - some of the confidence regions could even be empty! We have succeeded in deriving confidence regions that are intuitively appealing, are applicable to finite samples and are free of the drawbacks associated with the existing procedures. Furthermore, when applied to actual data, our confidence regions turned out to be much more satisfactory compared to those based on the existing procedures. The examples and simulation results indicate that we finally have a satisfactory solution to this problem, applicable to small samples.

The problem addressed in paper #2 is that of combining independent tests regarding a common parameter that occurs in the mean of several multivariate linear models. This is a problem in meta-analysis and has important applications in the social and behavioral sciences. This area of research also has applications in industrial robotics, commercial air traffic control and military operations where sensory data from multiple sensors are very often available and the problem is to combine diverse sets of sensed information. When the data can be modelled using a linear model, our results in paper #2 achieve this for hypotheses testing problems. Our results in this regard generalize the corresponding univariate results in the papers by Zhou and Mathew (1993, Journal of the American Statistical Association) and Mathew, Sinha and Zhou (1993, Journal of the American Statistical Association).

Satterthwaite's approximation of the distribution of a linear combination of independent means squares is a well known procedure in the analysis of univariate balanced mixed models. The multivariate analogue of this approximation is not so well known. This approximation applies to a nonnegative linear combination of independent Wishart matrices. In the article #3, we study the closeness of this approximation. A necessary and sufficient condition for the approximation to be exact is presented. A multivariate test statistic is then developed to detect any significant departure from this condition.

In the article #5, we establish a matrix inequality that provides an upper bound for a quadratic form that involves the difference between two linear unbiased estimators of the same linear parametric function in a general linear model. Various special cases of the inequality are discussed. Certain inequalities that arise in the problem of outlier detection and the problem of combining information from two independent linear models come out as special cases.

The work that is being completed in paper #7 deals with the nonnegative estimation of variance components in some time series models when measurement noise is present. The standard estimator in the time series literature is the maximum likelihood estimator, which can be computationally intense. We cast the time series model in direct correspondence with a mixed effects model and then use the results in the papers by Kelly and Mathew (Journal of the Royal Statistical Society, 1993, and Technometrics, 1994) in order to arrive at nonnegative estimators of variance components that have explicit analytic expressions, and hence are computationally easy. Furthermore, simulation results show that our estimators exhibit excellent performance (in terms of mean squared error) compared to the maximum likelihood estimator.

As a concluding remark, I would like to add that the most important research accomplishment among the above is the work on calibration, as mentioned at the beginning of this report. The results that I obtained in this context (in papers #1, #4 and #6) are exclusively on the construction of confidence regions. Papers #4 and #6 are jointly with a graduate student, W. Zha, also supported on the AFOSR grant, and will form part of her Ph. D. dissertation. (She will be graduating in the summer of 1995). Along with another graduate student, I am currently working on some point estimation problems that arise in multivariate calibration.